

On Partially Additive Kleene Algebras*

Riccardo Pucella
Cornell University
Ithaca, NY 14853 USA
riccardo@cs.cornell.edu

Abstract

We define the notion of a partially additive Kleene algebra, which is a Kleene algebra where the $+$ operation need only be partially defined. These structures formalize a number of examples that cannot be handled directly by Kleene algebras. We relate partially additive Kleene algebras to existing algebraic structures, by exhibiting categorical connections with Kleene algebras, partially additive categories, and closed semirings.

1 Introduction

Kleene algebras are algebraic structures that capture a natural form of iteration. Formally, a Kleene algebra is an idempotent semiring with a unary $*$ operation; it is this operation that is axiomatized using properties one expects from iteration. Kleene algebras are well-understood, having been studied since at least the work of Conway [1971], and provide a foundation for many program logics. The goal of this paper is to extend the theory of Kleene algebras to capture situations where it is possible to define a reasonable definition of iteration without there being a complete Kleene algebra structure. One example, studied by Kozen [1998], is the set of nonsquare matrices over a Kleene algebra, which does not have a multiplication operation defined for all pairs of matrices. This example can be studied by a form of Kleene algebra with a partially defined multiplication operation. We are interested in situations that require a different form of partiality, namely, where the addition operation is partial. An example where a partially defined addition operation is needed is given (essentially) by the set of partial functions on a set S , that is, the partial functions $S \rightarrow S$. We can define multiplication of two partial functions by composition. Similarly, we can define the sum of two partial functions f and g as $f \cup g$ (when considered as relations between arguments and results) provided that the values of f and g agree on their domain of definition; thus, addition of partial functions is only partially defined. Finally, we can define f^* as a form of reflexive transitive closure of f . Of course, partial functions can be studied using a number of approaches. This simply shows that we can bring them (and other examples) under the umbrella of a well-studied algebraic theory such as that of Kleene algebras.

Addition and multiplication having different properties in Kleene algebras, a theory for a partially defined addition operation is different than the theory for a partially defined multiplication

*A preliminary version of this paper appears in the proceedings of the 8th International Conference on Relational Methods in Computer Science (RelMiCS 8).

operation. This means that the development of Kozen [1998] for a partially defined multiplication is not directly applicable. Of course, the need for a partially defined addition has long been recognized, and a common way to deal with this issue is to consider a Kleene algebra with a special element $\star$ to represent the value “undefined”. When the sum of two elements is meant to be undefined, it is assigned the value $\star$, and applying any operation to a $\star$ argument results in $\star$. This is essentially the approach used by Kozen [2003] in a recent series of papers to perform dataflow analyses using Kleene algebras. While this method has the definite advantage of being simple and does not require any extension to the theory of Kleene algebras, it may not be always adequate. Among other things, it is not a *universal* way to add an undefined value to a Kleene algebra. For instance, we could imagine adding a number of undefined values, one per pair of elements whose sum is undefined, and defining addition appropriately for those values. There are properties of Kleene algebras extended with many undefined values that do not hold when a single undefined value is used, especially as it pertains to homomorphisms between Kleene algebras. What we want, for the purpose of developing a theory of Kleene algebras with a partially defined addition, is to identify the “minimal” way in which undefined values can be added, for an appropriate sense of minimality. It is not clear, a priori, that there is such a minimal extension.

To establish that such a minimal extension indeed exists, we proceed as follows. We define a primitive notion of *partially additive* Kleene algebra, which is a Kleene algebra where the $+$ operation is partial, rather than total. We establish the basic result that every star-continuous partially additive Kleene algebra (a class of algebras where the $*$ operation satisfies some additional and natural properties) can be embedded in a total Kleene algebra, and that this embedding is universal. Thus, there is a universal way to complete (star-continuous) partially additive Kleene algebras. This universal completion yields the minimal extension alluded to above. The remainder of the paper describes the relationship between partially additive Kleene algebras and other algebraic structures, with an eye towards showing that such Kleene algebras fit naturally within existing structures. First, we relate partially additive Kleene algebras with another structure that has been used to reason algebraically about iteration in the context of categorical semantics of programming languages, namely partially additive idempotent semirings and their generalization as partially additive categories [Manes and Arbib 1986]. We exhibit an adjunction between the category of star-continuous partially additive Kleene algebras and the category of partially additive idempotent semirings. Furthermore, we show that star-continuous partially additive Kleene algebras can be completed into closed semirings [Aho, Hopcroft, and Ullman 1975] in two distinct ways that nevertheless yield isomorphic closed semirings. This is done by exhibit two adjunctions between the category of star-continuous partially additive Kleene algebras and the category of closed semirings.

We assume a working knowledge of category theory, at the level of the first five chapters of MacLane [1971].

2 Partially Additive Kleene Algebras

We take as a starting point the definition of a Kleene algebra due to Kozen [1994], who gives a finitary axiomatization using implicational equations.

A *partially additive Kleene algebra* is a set $\mathcal{K}$ equipped with a partial binary operation $+$, a total binary operation $\cdot$, a total unary operation $*$, and constants 0 and 1, satisfying a number of axioms.

We say the pair (a, b) is *summable* (written $a \downarrow b$) if $a + b$ is defined. The axioms are as follows:

$$\text{if } x \downarrow y, (x + y) \downarrow z, \text{ and } y \downarrow z \text{ then } x \downarrow (y + z) \text{ and } (x + y) + z = x + (y + z) \quad (1)$$

$$\text{if } x \downarrow y \text{ then } y \downarrow x \text{ and } x + y = y + x \quad (2)$$

$$x \downarrow 0 \text{ and } x + 0 = x \quad (3)$$

$$x \downarrow x \text{ and } x + x = x \quad (4)$$

$$x \cdot (y \cdot z) = (x \cdot y) \cdot z \quad (5)$$

$$1 \cdot x = x \quad (6)$$

$$x \cdot 1 = x \quad (7)$$

$$\text{if } x \downarrow y \text{ then } (z \cdot x) \downarrow (z \cdot y) \text{ and } z \cdot (x + y) = z \cdot x + z \cdot y \quad (8)$$

$$\text{if } x \downarrow y \text{ then } (x \cdot z) \downarrow (y \cdot z) \text{ and } (x + y) \cdot z = x \cdot z + y \cdot z \quad (9)$$

$$0 \cdot x = 0 \quad (10)$$

$$x \cdot 0 = 0 \quad (11)$$

$$1 \leq x^* \quad (12)$$

$$x \cdot x^* \leq x^* \quad (13)$$

$$x^* \cdot x \leq x^* \quad (14)$$

$$\text{if } a \cdot x \leq x \text{ then } a^* \cdot x \leq x \quad (15)$$

$$\text{if } x \cdot a \leq x \text{ then } x \cdot a^* \leq x \quad (16)$$

As usual, every variable appearing in an axiom is implicitly understood as universally quantified. The relation $\leq$ in axioms (12)–(16) is the natural partial order on $\mathcal{K}$ induced by $+$: $a \leq b$ if and only if $a \downarrow b$ and $a + b = b$. It is straightforward to check that $\leq$ is symmetric (since $a \downarrow 0$), transitive (by the associativity rules for summability), and antisymmetric (using the idempotence of $+$). It is easy to check that $+$ captures a form of least upperbound: if $a \leq c$, $b \leq c$, and $a \downarrow b$, then $a + b \leq c$ (using associativity rules for summability). We often drop the $\cdot$ symbol for multiplication, simply writing ab for $a \cdot b$. For b an element of a partially additive Kleene algebra, define b^0 to be 1, and b^{n+1} to be bb^n .

Axioms (1)–(4) say that $(\mathcal{K}, +, 0)$ is an idempotent commutative monoid, except that the $+$ operation is partial, while axioms (5)–(9) say that $(\mathcal{K}, \cdot, 1)$ is a monoid, with the $\cdot$ operation distributing over $+$ whenever the required sums are defined. With axioms (10)–(11), this makes $(\mathcal{K}, +, \cdot, 0, 1)$ an idempotent semiring with a partially defined $+$ operation. Axioms (14)–(17) capture the properties of the Kleene star operation of formal language theory, taking into account the partiality of $+$. Most of the $\leq$ -related properties that hold in Kleene algebra also hold here, even taking into account partial additivity (since $a \leq b$ requires that $a \downarrow b$). For instance, we can check that for all $n \geq 0$, $b^n \leq b^*$ (using induction from (12)). Similarly, $a^*a^* = a^*$, $(a^*)^* = a^*$, and if $a \leq b$, then $a^* \leq b^*$. Rather than listing all the properties of interest here, we refer to Kozen [1994].

Note that only $+$ is allowed to be partial in a partially additive Kleene algebra. This lets us relate our results to partially additive semirings in the next section, and also captures the kind of examples we want to capture. Allowing $\cdot$ to be partial is essentially an orthogonal feature. A form of Kleene algebra where the $\cdot$ operation is partial is studied by Kozen [1998] under the name *typed Kleene algebra*, where a type system is used to control the partiality of multiplication. The definitions and results of this paper should carry over in a straightforward way to that setting. (We do not introduce

a type system in this paper to control the partiality of addition, since it would make the framework too restrictive for the examples we want to capture.)

Example 2.1. Every total Kleene algebra is a partially additive Kleene algebra where the $+$ operation is in fact total. $\square$

Example 2.2. Any commutative monoid can be turned into a partially additive Kleene algebra with a maximally undefined operation $+$. Let $(\mathcal{M}, \cdot, 1)$ be a commutative monoid. Let $\mathcal{M}' = \mathcal{M} \cup \{0, \star\}$, where 0 and $\star$ are new elements, and extend $\cdot$ to $\mathcal{M}'$ by taking $0 \cdot a = a \cdot 0 = 0$ for all $a \in \mathcal{M}'$, and $\star \cdot b = b \cdot \star = \star$ for all $b \in \mathcal{M}'$, $b \neq 0$. Define $0^* = 0$, and $b^* = \star$ (if $b \neq 0$). Let $a + b$ be defined only if $a = b$, in which case $a + b = a$, if $b = 0$, in which case $a + b = a$, or if $b = \star$, in which case $a + b = b$. It is easy to check that $(\mathcal{M}', +, \cdot, *, 0, 1)$ forms a partially additive Kleene algebra. $\square$

Example 2.3. Let Σ be a set of states. In much the same way that relations are the basic example of Kleene algebras, the set of partial functions from Σ to Σ is a basic example of a partially additive Kleene algebra, where we interpret f^* as repeatedly applying f until the state is in a distinguished set $\Omega \subseteq \Sigma$. Unfortunately, the example is not quite as simple as in the total case. Intuitively, we need the partial functions to record some information about how they have been composed. Therefore, we look at strings over Σ , where we can interpret a string $\sigma_1 \dots \sigma_n$ as mapping σ_1 to σ_n (via $\sigma_2, \dots, \sigma_{n-1}$). Let ϵ be the empty string. For a nonempty string $s = \sigma_1 \dots \sigma_n$, let s_{ini} , the initial state of s , be σ_1 , and s_{fin} , the final state of s , be σ_n . Given two strings s and s' , say that s is a generalized prefix of s' if $s' = s_1 t_1 s_2 t_2 \dots s_n t_n$ and $s = s_1 s_2 \dots s_n$. The fusion product $s \otimes s'$ of two strings is defined to be ϵ if either s or s' is ϵ , or if $s_{\text{fin}} \neq s'_{\text{ini}}$. Otherwise, if $s = \sigma_1 \dots \sigma_m$ and $s' = \sigma'_1 \dots \sigma'_n$, then $s \otimes s' = \sigma_1 \dots \sigma_m \sigma'_2 \dots \sigma'_n$. Let $\Omega \subseteq \Sigma$ be a fixed subset of Σ . A set $A \subseteq \Sigma^*$ of strings is *functional* if (1) for all $s = \sigma_1 \dots \sigma_n \in A$, if $\sigma_i \in \Omega$, then $i = n$ (only the last state of a string is in Ω), (2) for all $\sigma \in \Omega$, there is a string $s \in A$ with $s_{\text{ini}} = \sigma$, and (3) for all $\sigma \in \Sigma$ and all maximal strings $s, s' \in A$, $s_{\text{fin}} = s'_{\text{fin}}$, where a string s is maximal in A if it is not a generalized prefix of any other string in A . A set $A \subseteq \Sigma^*$ is *sparsely functional* if it is functional, and moreover no string in A is a generalized prefix of any other string in A . Let $\mathcal{F}$ be the set of sparsely functional subsets of Σ^* , along with the empty set. Given A a functional set, let $\lceil A \rceil$ be the largest set $A' \subseteq A$ that is sparsely functional. The sparsely functional subsets of Σ^* form a partially additive Kleene algebra $(\mathcal{F}, +, \cdot, *, 0, 1)$ where we take $A \downarrow B$ if $A \cup B$ is functional, in which case $A + B = \lceil A \cup B \rceil$, $A \cdot B = \lceil \{s \otimes s' \mid s \in A, s' \in B\} \rceil$, and $A^* = \lceil \Sigma \cup \{s \mid \exists n (s \in A^n, s_{\text{fin}} \in \Omega) \} \rceil$, where $A^0 = \Omega$, and $A^{n+1} = A \cdot A^n$. Thus, if we interpret A as a partial function from the initial states of the strings in A to the final states of these strings, then $A + B$ represents the union of partial functions, $A \cdot B$ represents the composition of partial functions, and A^* represents the iterated application of A until an element of Ω is reached. The constant 0 is taken to be the empty set, and the constant 1 is taken to be Σ . $\square$

A homomorphism of partially additive Kleene algebras is a function $f : \mathcal{K} \rightarrow \mathcal{K}'$ such that if $a \downarrow b$ in $\mathcal{K}$, then $f(a) \downarrow f(b)$ in $\mathcal{K}'$ and $f(a + b) = f(a) + f(b)$, as well as $f(a \cdot b) = f(a) \cdot f(b)$, $f(a^*) = f(a)^*$, $f(0) = 0$, and $f(1) = 1$. Partially additive Kleene algebras with homomorphisms form a category **PKA**. It follows from Example 2.1 that the category **KA** of Kleene algebra forms a full subcategory of **PKA**, via the inclusion functor $P : \mathbf{KA} \rightarrow \mathbf{PKA}$.

Note that every partially additive Kleene algebra embeds in a total Kleene algebra in a straightforward way, by adding a single element $\star$. More precisely, let $\mathcal{K}$ be a partially additive Kleene

algebra, and consider the Kleene algebra $\mathcal{K} \cup \{\star\}$ with the same operations, extended so that if a and b are not summable in $\mathcal{K}$, then $a + b = \star$ in $\mathcal{K} \cup \{\star\}$, with $a + \star = \star$, $0 \cdot \star = \star \cdot 0 = 0$, $a \cdot \star = \star \cdot a = \star$, and $\star^\star = \star$. It is easy to check that $\mathcal{K} \cup \{\star\}$ is a Kleene algebra with a total operation $+$. Unfortunately, this embedding is not universal. Intuitively, this construction does not embed a partially additive Kleene algebra in the “closest” Kleene algebra that contains it. We will see a universal embedding in the next section.

3 Star-Continuity

A particularly interesting class of partially additive Kleene algebras is one where the Kleene star operator is related to suprema with respect to $\leq$. A partially additive Kleene algebra is *star-continuous* if $ab^\star c = \sum_{n < \omega} ab^n c$, where $\sum$ is the supremum with respect to the order $\leq$. This equation is equivalent to the infinite implication

$$\text{if } ab^n c \downarrow w \text{ for all } n \text{ and } \bigwedge_{n < \omega} ab^n c \leq w \text{ then } ab^\star c \leq w. \quad (17)$$

Equation (17) does not always hold in partially additive Kleene algebras, simply because not all Kleene algebras are star-continuous [Kozen 1990]. However, most natural examples of partially additive Kleene algebra are in fact star-continuous. In particular, Examples 2.2 and 2.3 of the last section are star-continuous.

3.1 Construction

Star-continuous partially additive Kleene algebras form a full subcategory $\mathbf{PKA}^\star$ of $\mathbf{PKA}$. We now show that the inclusion functor $K : \mathbf{PKA}^\star \rightarrow \mathbf{PKA}$ witnessing the fact that $\mathbf{PKA}^\star$ is a subcategory has a left adjoint S that universally maps a partially additive Kleene algebra to the “closest” star-continuous partially additive Kleene algebra. The idea is to take a partially additive Kleene algebra and impose equation (17) upon it. (The idea is similar to the Abelianization of groups, which universally maps a group to an Abelian group.) Note that there is nothing specific to partially additive Kleene algebra here; this construction equally well maps a Kleene algebra to the closest star-continuous Kleene algebra, and forms a left adjoint to the appropriate inclusion functor.

Constructing the left adjoint to K can be done as follows. Let $\mathcal{K}$ be a partially additive Kleene algebra. Define $\equiv$ to be the least congruence relation on $\mathcal{K}$ such that for all a, b, c, w ,

$$\text{if } ab^n c + w \equiv w \text{ (for all } n \geq 0) \text{ then } ab^\star c + w \equiv w. \quad (18)$$

(In the presence of partiality, congruence for $+$ says that if $x \downarrow z$, $y \downarrow z$, and $x \equiv y$, then $x + z \equiv y + z$.) Let $[x]_\equiv = \{y \mid x \equiv y\}$ be the equivalence class of x under congruence $\equiv$. Let $\mathcal{K}/\equiv$ be the set of equivalence classes of $\equiv$. If we extend the operations of $\mathcal{K}$ to equivalence classes in the standard way, it is easily seen that $\mathcal{K}/\equiv$ is a partially additive Kleene algebra that is moreover star-continuous. (Say $[x]_\equiv \downarrow [y]_\equiv$ if there exists $x' \equiv x$ and $y' \equiv y$ such that $x' \downarrow y'$, and set $[x]_\equiv + [y]_\equiv = [x' + y']_\equiv$.) If $f : \mathcal{K} \rightarrow \mathcal{K}'$ is a partially additive Kleene algebra homomorphism, we can lift it to a homomorphism $\hat{f} : \mathcal{K}/\equiv \rightarrow \mathcal{K}'/\equiv$ by taking $\hat{f}([x]_\equiv) = [f(x)]_\equiv$.

Lemma 3.1. *If f is a partially additive Kleene algebra homomorphism, then $\hat{f}$ is a partially additive Kleene algebra homomorphism.*

Proof. The main thing we need to check is that $\hat{f}$ is well-defined, that is, if $x \equiv y$, then $\hat{f}([x]_{\equiv}) = \hat{f}([y]_{\equiv})$, that is, $f(x) \equiv f(y)$. Let $h : \mathcal{K}' \rightarrow \mathcal{K}'/\equiv$ be the canonical surjective homomorphism mapping an element x of $\mathcal{K}'$ to the equivalence class $[x]_{\equiv}$ of $\mathcal{K}'/\equiv$. Recall that the kernel $\ker(g)$ of a homomorphism $g : \mathcal{K} \rightarrow \mathcal{K}'$ is the set of all pairs of elements (x, y) such that $g(x) = g(y)$. We show that $\equiv \subseteq \ker(h \circ f)$, so that $x \equiv y$ implies that $h(f(x)) = h(f(y))$, that is, $[f(x)]_{\equiv} = [f(y)]_{\equiv}$, as required. Recall that the kernel of a homomorphism is a congruence; this holds even in the presence of partiality for $+$: if $x \downarrow z, y \downarrow z$, and $(x, y) \in \ker f$, then $f(x+z) = f(x)+f(z) = f(y)+f(z) = f(y+z)$, so that $(x+z, y+z) \in \ker f$. Therefore, $\ker(h \circ f)$ is a congruence. It therefore suffices to show that $\ker(h \circ f)$ is closed under (17) to get $\equiv \subseteq \ker h \circ f$. Assume that for all n , $ab^n c + c \downarrow w$ and $(ab^n c + w, w) \in \ker(h \circ f)$. In other words, we have $h(f(ab^n c + w)) = h(f(w))$ for all n . Because $h \circ f$ is a homomorphism, we get $h(f(a))h(f(b))^n h(f(c)) + h(f(w)) = h(f(w))$ for all n . Since $\mathcal{K}'/\equiv$ is star-continuous, $h(f(a))h(f(b))^* h(f(c)) + h(f(w)) = h(f(w))$ holds, and therefore, $h(f(ab^* c + w)) = h(f(w))$; that is, $(ab^* c + w, w) \in \ker(h \circ f)$, as required.

Verifying that $\hat{f}$ is a partially additive Kleene algebra homomorphism is straightforward. For instance, if $[x]_{\equiv} \downarrow [y]_{\equiv}$, then $\hat{f}([x]_{\equiv} + [y]_{\equiv}) = \hat{f}([x+y]_{\equiv}) = [f(x+y)]_{\equiv} = [f(x) + f(y)]_{\equiv} = [f(x)]_{\equiv} + [f(y)]_{\equiv} = \hat{f}([x]_{\equiv}) + \hat{f}([y]_{\equiv})$. $\square$

Let $S : \mathbf{PKA} \rightarrow \mathbf{PKA}^*$ map $\mathcal{K}$ to $\mathcal{K}/\equiv$, and f to $\hat{f}$. It is easy to check that S is a functor.

Theorem 3.2. *The functor S is a left adjoint of the inclusion functor $K : \mathbf{PKA}^* \rightarrow \mathbf{PKA}$, via the adjunction map $\varphi : \text{Hom}(SK, \mathcal{K}') \rightarrow \text{Hom}(\mathcal{K}, K\mathcal{K}')$ given by $(\varphi f)(x) = f([x]_{\equiv})$.*

Proof. To show that S is a left adjoint to K , we need to show that the adjunction map φ is a natural isomorphism between $\text{Hom}(SK, \mathcal{K}')$ and $\text{Hom}(\mathcal{K}, K\mathcal{K}')$.

We first show that φ is an isomorphism. The inverse of φ is given by the map ψ defined by $(\psi g)([x]_{\equiv}) = g(x)$. We check that this function is well defined, that is, if $x \equiv y$, then $g(x) = g(y)$. The argument is similar to that in the proof of Lemma 3.1. Clearly, it suffices to show that $\equiv \subseteq \ker(g)$. Since the kernel of a homomorphism is a congruence, it suffices to show that $\ker(g)$ is closed under (17). Let $g : \mathcal{K} \rightarrow K\mathcal{K}'$, where $\mathcal{K}'$ is a star-continuous partially additive Kleene algebra. Assume that for all $n \geq 0$, $ab^n c \downarrow w$ and $(ab^n c + w, w) \in \ker(g)$, that is, $g(ab^n c + w) = g(w)$ for all $n \geq 0$. Since g is a homomorphism, we get $g(a)g(b)^n g(c) + g(w) = g(w)$ for all $n \geq 0$. Since $\mathcal{K}'$ is star-continuous (and K is the forgetful functor), we have $g(a)g(b)^* g(c) + g(w) = g(w)$, that is, $g(ab^* c + w) = g(w)$, and $(ab^* c + w, w) \in \ker(g)$, establishing the result. It is easy to see that (ψg) is a partially additive Kleene algebra homomorphism. To see that φ and ψ are inverses, first let $g : \mathcal{K} \rightarrow K\mathcal{K}'$, and note that

$$\begin{aligned} (\varphi \circ \psi)(g)(x) &= (\varphi(\psi(g)))(x) \\ &= (\psi(g))([x]_{\equiv}) \\ &= g(x). \end{aligned}$$

Second, let $f : SK \rightarrow \mathcal{K}'$, and note that

$$\begin{aligned} (\psi \circ \varphi)(f)([x]_{\equiv}) &= (\psi(\varphi(f)))([x]_{\equiv}) \\ &= (\varphi f)(x) \\ &= f([x]_{\equiv}). \end{aligned}$$

To establish the naturality of the adjunction map φ , we need to show the commutativity of two diagrams. First, we need to show that for all morphisms $h : \mathcal{K}' \rightarrow \mathcal{K}''$ in $\mathbf{PKA}^*$, the following diagram commutes:

$$\begin{array}{ccc} \mathrm{Hom}(S\mathcal{K}, \mathcal{K}') & \xrightarrow{\varphi} & \mathrm{Hom}(\mathcal{K}, K\mathcal{K}') \\ \mathrm{Hom}(S\mathcal{K}, h) \downarrow & & \downarrow \mathrm{Hom}(\mathcal{K}, Kh) \\ \mathrm{Hom}(S\mathcal{K}, \mathcal{K}'') & \xrightarrow{\varphi} & \mathrm{Hom}(\mathcal{K}, K\mathcal{K}'') \end{array}$$

where $\mathrm{Hom}(\mathcal{K}, -)$ is a functor that maps an object $\mathcal{K}'$ to the set $\mathrm{Hom}(\mathcal{K}, \mathcal{K}')$, and a morphism $h : \mathcal{K}' \rightarrow \mathcal{K}''$ to a morphism $\mathrm{Hom}(\mathcal{K}, h) : \mathrm{Hom}(\mathcal{K}, \mathcal{K}') \rightarrow \mathrm{Hom}(\mathcal{K}, \mathcal{K}'')$, given by $\mathrm{Hom}(\mathcal{K}, h)(f) = h \circ f$. To verify the commutativity of the diagram, let $h : \mathcal{K}' \rightarrow \mathcal{K}''$, let $f : S\mathcal{K} \rightarrow \mathcal{K}'$, and let $x \in \mathcal{K}$:

$$\begin{aligned} (\mathrm{Hom}(\mathcal{K}, Kh) \circ \varphi)(f)(x) &= (\mathrm{Hom}(\mathcal{K}, Kh)(\varphi(f)))(x) \\ &= ((Kh) \circ (\varphi f))(x) \\ &= ((Kh)((\varphi f)(x))) \\ &= h(f([x]_{\equiv})) \end{aligned}$$

(since $Kh = h$) and

$$\begin{aligned} (\varphi \circ \mathrm{Hom}(S\mathcal{K}, h))(f)(x) &= (\varphi(\mathrm{Hom}(S\mathcal{K}, h)(f)))(x) \\ &= (\varphi(h \circ f))(x) \\ &= (h \circ f)([x]_{\equiv}) \\ &= h(f([x]_{\equiv})). \end{aligned}$$

Second, we need to show that for all morphisms $h : \mathcal{K}'' \rightarrow \mathcal{K}'$ in $\mathbf{PKA}$, the following diagram commutes:

$$\begin{array}{ccc} \mathrm{Hom}(S\mathcal{K}', \mathcal{K}) & \xrightarrow{\varphi} & \mathrm{Hom}(\mathcal{K}', K\mathcal{K}) \\ \mathrm{Hom}(Sh, \mathcal{K}) \downarrow & & \downarrow \mathrm{Hom}(h, K\mathcal{K}) \\ \mathrm{Hom}(S\mathcal{K}'', \mathcal{K}) & \xrightarrow{\varphi} & \mathrm{Hom}(\mathcal{K}'', K\mathcal{K}) \end{array}$$

where $\mathrm{Hom}(-, \mathcal{K})$ is a functor that maps an object $\mathcal{K}'$ to the set $\mathrm{Hom}(\mathcal{K}', \mathcal{K})$, and a morphism $h : \mathcal{K}'' \rightarrow \mathcal{K}'$ to a morphism $\mathrm{Hom}(h, \mathcal{K}) : \mathrm{Hom}(\mathcal{K}', \mathcal{K}) \rightarrow \mathrm{Hom}(\mathcal{K}'', \mathcal{K})$, given by $\mathrm{Hom}(h, \mathcal{K})(f) = f \circ h$. To verify the commutativity of the diagram, let $h : \mathcal{K}'' \rightarrow \mathcal{K}'$, let $f : S\mathcal{K}' \rightarrow \mathcal{K}$, and let $x \in \mathcal{K}''$:

$$\begin{aligned} (\mathrm{Hom}(h, K\mathcal{K}) \circ \varphi)(f)(x) &= (\mathrm{Hom}(h, K\mathcal{K})(\varphi f))(x) \\ &= ((\varphi f) \circ h)(x) \\ &= (\varphi f)(h(x)) \\ &= f([h(x)]_{\equiv}) \end{aligned}$$

and

$$\begin{aligned}
(\varphi \circ \text{Hom}(Sh, \mathcal{K}))(f)(x) &= (\varphi(\text{Hom}(Sh, \mathcal{K})(f)))(x) \\
&= (\varphi(f \circ (Sh)))(x) \\
&= (\varphi(f \circ \hat{h}))(x) \\
&= (f \circ \hat{h})([x]_{\equiv}) \\
&= f(\hat{h}([x]_{\equiv})) \\
&= f([h(x)]_{\equiv}).
\end{aligned}$$

□

3.2 Completion

More interesting is the relationship between star-continuous Kleene algebras and star-continuous partially additive Kleene algebras. In one direction, every star-continuous Kleene algebra is a star-continuous partially additive Kleene algebra. Therefore, the inclusion functor P , when restricted to the full subcategory $\mathbf{KA}^*$ of star-continuous Kleene algebra, gives rise to an inclusion functor (also named P) from $\mathbf{KA}^*$ to $\mathbf{PKA}^*$. In the other direction, every star-continuous partially additive Kleene algebra $\mathcal{K}$ can be *completed* to a star-continuous Kleene algebra $T\mathcal{K}$ in a universal way. Formally, we show how to define a functor T from the category $\mathbf{PKA}^*$ to the category $\mathbf{KA}^*$ that is the left adjoint to the inclusion functor P .

The construction is a form of ideal completion, and is based on a construction of Conway [1971], later formalized by Kozen [1990]. We follow that treatment rather closely. A *star-ideal* A in a star-continuous partially additive Kleene algebra $\mathcal{K}$ is a subset $A \subseteq \mathcal{K}$ satisfying:

- (1) A is nonempty;
- (2) A is closed under $+$ (for summable pairs);
- (3) A is closed downward under $\leq$;
- (4) if $ab^n c$ is in A for all n , then $ab^* c$ is in A .

The *star-ideal generated by* A , denoted $\langle A \rangle$, is the smallest ideal containing A . It is equivalently defined as the intersection of all the star-ideals containing A . If $A = \{a_1, \dots, a_k\}$, we often write $\langle a_1, \dots, a_k \rangle$ for $\langle A \rangle$. In particular, if $A = \{a\}$, then we write $\langle a \rangle$ for $\langle A \rangle$. It is easy to check that $\langle a \rangle = \{x \mid x \leq a\}$. An ideal I is *finitely generated* if $I = \langle A \rangle$ for some finite set A , and *countably generated* if $I = \langle A \rangle$ for some countable set A . We can construct the star-ideal generated by A directly, using a transfinite argument. This is a useful technique for proving results, as we shall see.

Lemma 3.3. *Let τ be the map $\tau(A) = A \oplus A \cup A^{\leq} \cup A^{*}$, where*

$$\begin{aligned}
A \oplus B &= \{a + b \mid a \in A, b \in B, a, b \text{ summable}\} \\
A^{\leq} &= \{y \mid \exists x \in A. y \leq x\} \\
A^{*} &= \{ab^* c \mid ab^n c \in A, \text{ for all } n \geq 0\},
\end{aligned}$$

and define the transfinite sequence

$$\begin{aligned}\tau^0(A) &= A \\ \tau^{\alpha+1}(A) &= \tau(\tau^\alpha(A)) \\ \tau^\lambda(A) &= \bigcup_{\alpha < \lambda} \tau^\alpha(A) \quad \lambda \text{ a limit ordinal.}\end{aligned}$$

For all $A \subseteq \mathcal{K}$, there exists an ordinal κ such that $\tau^\kappa(A) = \tau^{\kappa+1}(A)$, and $\tau^\kappa = \langle A \rangle$.

Proof. It is easy to check that $A \subseteq \tau(A)$, and that τ is monotone, that is, $\tau(A) \subseteq \tau(B)$ whenever $A \subseteq B$. The result then follows immediately by the Knaster-Tarski theorem [Tarski 1955]. $\square$

Take $T\mathcal{K}$ to be the set of all finitely generated star-ideals of $\mathcal{K}$. We can impose a star-continuous Kleene algebra structure on $T\mathcal{K}$ as follows. First, define the following operation $\odot$ on subsets of $\mathcal{K}$:

$$A \odot B = \{a \cdot b \mid a \in A, b \in B\}.$$

Lemma 3.4. *The following properties hold:*

- (1) $\langle A \odot B \rangle = \langle \langle A \rangle \odot B \rangle = \langle A \odot \langle B \rangle \rangle$;
- (2) $\langle A \cup B \rangle = \langle \langle A \rangle \cup B \rangle = \langle A \cup \langle B \rangle \rangle$.

Proof. We prove part (2) here; part (1) is essentially Lemma 3 in Kozen [1990], and is proved similarly. Since $\cup$ is commutative, it is sufficient to show that $\langle A \cup B \rangle = \langle \langle A \rangle \cup B \rangle$. Since $A \subseteq \langle A \rangle \subseteq \langle \langle A \rangle \cup B \rangle$ and $B \subseteq \langle \langle A \rangle \cup B \rangle$, we have $\langle A \cup B \rangle \subseteq \langle \langle A \rangle \cup B \rangle$. We show the reverse inclusion $\langle \langle A \rangle \cup B \rangle \subseteq \langle A \cup B \rangle$ by transfinite induction, via the characterization of star-ideals in Lemma 3.3. More precisely, we show that for all ordinals α , $\tau^\alpha(\langle \langle A \rangle \cup B \rangle) \subseteq \langle A \cup B \rangle$. It is easy to see that since $A \subseteq \langle A \cup B \rangle$, $\langle A \rangle \subseteq \langle A \cup B \rangle$, and similarly, $B \subseteq \langle A \cup B \rangle$, so that $\tau^0(\langle \langle A \rangle \cup B \rangle) = \langle A \rangle \cup B \subseteq \langle A \cup B \rangle$. For the inductive case,

$$\begin{aligned}\tau^{\alpha+1}(\langle \langle A \rangle \cup B \rangle) &= \tau(\tau^\alpha(\langle \langle A \rangle \cup B \rangle)) \\ &\subseteq \tau(\langle A \cup B \rangle) \quad \text{by monotonicity of } \tau \\ &= \langle A \cup B \rangle \quad \text{by property of star-ideals.}\end{aligned}$$

For limit ordinals λ ,

$$\begin{aligned}\tau^\lambda(\langle \langle A \rangle \cup B \rangle) &= \bigcup_{\alpha < \lambda} \tau^\alpha(\langle \langle A \rangle \cup B \rangle) \\ &\subseteq \bigcup_{\alpha < \lambda} \langle A \cup B \rangle \\ &= \langle A \cup B \rangle.\end{aligned}$$

Thus, $\langle \langle A \rangle \cup B \rangle = \tau^*(\langle \langle A \rangle \cup B \rangle) \subseteq \langle A \cup B \rangle$, as required. $\square$

Clearly, Lemma 3.4 extends to arbitrary finite products and unions. If I and J are finitely generated star-ideals, generated respectively by $A = \{a_1, \dots, a_k\}$ and B , define

$$\begin{aligned} I + J &= \langle A \cup B \rangle \\ I \cdot J &= \langle A \odot B \rangle \\ I^* &= \langle (a_1^* \dots a_k^*)^* \rangle \\ 1 &= \langle 1 \rangle \\ 0 &= \langle 0 \rangle. \end{aligned}$$

Note that the operation $+$ on star-ideals is total. We need to check that these operations are well-defined; there can be many different generators for a star-ideal, and the above operations need to give the same result no matter what generators we take.

Lemma 3.5. *If $I = \langle A_1 \rangle = \langle A_2 \rangle$ and $J = \langle B_1 \rangle = \langle B_2 \rangle$, then:*

- (1) $\langle A_1 \cup B_1 \rangle = \langle A_2 \cup B_2 \rangle$;
- (2) $\langle A_1 \odot B_1 \rangle = \langle A_2 \odot B_2 \rangle$;
- (3) *If $A_1 = \langle a_1, \dots, a_k \rangle$ and $B_1 = \langle b_1, \dots, b_l \rangle$, then $\langle (a_1^* \dots a_k^*)^* \rangle = \langle (b_1^* \dots b_l^*)^* \rangle$.*

Proof. For part (1), by Lemma 3.4(1), we have $\langle A_1 \cup B_1 \rangle = \langle \langle A_1 \rangle \cup \langle B_1 \rangle \rangle = \langle \langle A_2 \rangle \cup \langle B_2 \rangle \rangle = \langle A_2 \cup B_2 \rangle$.

For part (2), by Lemma 3.4(2), we have $\langle A_1 \odot B_1 \rangle = \langle \langle A_1 \rangle \odot \langle B_1 \rangle \rangle = \langle \langle A_2 \rangle \odot \langle B_2 \rangle \rangle = \langle A_2 \odot B_2 \rangle$.

For part (3), we first establish that for all i , $a_i \leq (b_1^* \dots b_l^*)^*$. This follows easily from the fact that $a_i \in \langle (a_1^* \dots a_k^*)^* \rangle$, and thus that $a_i \in \langle (b_1^* \dots b_l^*)^* \rangle$, and by principality, $a_i \leq (b_1^* \dots b_l^*)^*$. The result now follows almost immediately. For each i , by monotonicity of $*$, $a_i^* \leq ((b_1^* \dots b_l^*)^*)^* = (b_1^* \dots b_l^*)^*$. Thus, $a_1^* \dots a_k^* \leq (b_1^* \dots b_l^*)^* \dots (b_1^* \dots b_l^*)^* \leq (b_1^* \dots b_l^*)^*$. By monotonicity, $(a_1^* \dots a_k^*)^* \leq ((b_1^* \dots b_l^*)^*)^* = (b_1^* \dots b_l^*)^*$. Therefore, $\langle (a_1^* \dots a_k^*)^* \rangle \subseteq \langle (b_1^* \dots b_l^*)^* \rangle$. A symmetric argument gives us the reverse inclusion, establishing equality. $\square$

We can check that $\leq$ for star-ideals is just subset inclusion.

Lemma 3.6. *If $\mathcal{K}$ is a star-continuous partially additive Kleene algebra, then TK is a star-continuous Kleene algebra.*

Proof. Most of axioms of Kleene algebra (given by Kozen [1994]) are straightforward to verify. We give a proof for the nontrivial ones: $1 + I \cdot I^* \leq I^*$, $J + I \cdot K \leq K$ implies $I^* \cdot J \leq K$, and the star-continuity condition.

To show that $1 + I \cdot I^* \leq I^*$, it suffices to show that $1 \leq I^*$ and $I \cdot I^* \leq I^*$. Assume $I = \langle A \rangle$, where $A = \{a_1, \dots, a_k\}$. First, since $\mathcal{K}$ is a partially additive Kleene algebra, $1 \leq (a_1^* \dots a_k^*)^*$, and so $1 \in \langle (a_1^* \dots a_k^*)^* \rangle$, and $\langle 1 \rangle \subseteq \langle (a_1^* \dots a_k^*)^* \rangle$, as required. Second, pick an arbitrary $a \in A$. Clearly, $a \leq a_1^* \dots a_k^*$, so $a(a_1^* \dots a_k^*)^* \leq a_1^* \dots a_k^*(a_1^* \dots a_k^*)^* \leq (a_1^* \dots a_k^*)^*$. Therefore, by closure of I^* , $a(a_1^* \dots a_k^*)^* \in I^*$, for any $a \in A$. Thus, $A \odot \{(a_1^* \dots a_k^*)^*\} \subseteq I^*$, and $\langle A \odot \{(a_1^* \dots a_k^*)^*\} \rangle \subseteq I^*$. But this is just $I \cdot I^* \leq I^*$, as required.

Next, we want to show that if $J + I \cdot K \leq K$, then $I^* \cdot J \leq K$. Assume $I = \langle A \rangle$ (with $A = \{a_1, \dots, a_k\}$), $J = \langle B \rangle$, and $K = \langle C \rangle$. Furthermore, assume $J + I \cdot K \leq K$, that is, $\langle B \cup (A \odot C) \rangle \subseteq \langle C \rangle$. By Lemma 3.4, this is equivalent to

$$\langle \langle B \rangle \cup (\langle A \rangle \odot \langle C \rangle) \rangle \subseteq \langle C \rangle. \quad (19)$$

We want to show that $\langle \{(a_1^* \cdots a_k^*)^*\} \odot B \rangle \subseteq \langle C \rangle$. It is sufficient to show that for all $b \in B$, $(a_1^* \cdots a_k^*)^* b \in \langle C \rangle$. We know $b \in \langle B \rangle$, thus by (19), $b \in \langle C \rangle$. since $a_k \in \langle A \rangle$, $a_k b \in \langle A \rangle \odot \langle C \rangle$, and by (19), $a_k b \in \langle C \rangle$. A straightforward induction shows that for all n , $a_k^n b \in \langle C \rangle$. Therefore, by the closure properties of $\langle C \rangle$, $a_k^* b \in \langle C \rangle$. A similarly argument with $a_k^* b$ instead of b and a_{k-1} instead of a_k shows that $a_{k-1}^* a_k^* b \in \langle C \rangle$. A straightforward induction shows that $a_1^* \cdots a_k^* b \in \langle C \rangle$. Repeating the whole argument with $a_1^* \cdots a_k^* b$ instead of b yields $(a_1^* \cdots a_k^*)^2 b \in \langle C \rangle$. A straightforward induction shows that for all n , $(a_1^* \cdots a_k^*)^n b \in \langle C \rangle$. By the closure properties of $\langle C \rangle$, $(a_1^* \cdots a_k^*)^* b \in \langle C \rangle$, as required.

For the star-continuity condition, we want to show that if $I \cdot J^n \cdot K \leq L$ for all n , then $I \cdot J^* \cdot K \leq L$. Assume $I = \langle A \rangle$, $J = \langle B \rangle$ (where $B = \{b_1, \dots, b_k\}$), and $K = \langle C \rangle$. Furthermore, assume $I \cdot J^n \cdot K \leq L$ for all n , that is, $\langle A \odot B^n \odot C \rangle \subseteq L$, where $B^n = B \odot \cdots \odot B$ (n times). By Lemma 3.4, this is equivalent to

$$\langle \langle A \rangle \odot \langle B \rangle^n \odot \langle C \rangle \rangle \subseteq L. \quad (20)$$

It is sufficient to show that $A \odot \{(b_1^* \cdots b_k^*)^*\} \odot C \subseteq L$. In other words, for all $a \in A$ and $c \in C$, $a(b_1^* \cdots b_k^*)^* c \in L$. By the closure properties of L , it is sufficient to show that for all $a \in A$, all $c \in C$, and all $m \geq 0$, $a(b_1^* \cdots b_k^*)^m c \in L$. For $m = 0$, the result follows trivially from (20) (taking $n = 0$). For $m > 0$, by (20), for all $n_1^1, \dots, n_k^1, \dots, n_1^m, \dots, n_k^m$, we have $a(b_1^{n_1^1} \cdots b_k^{n_k^1}) \cdots (b_1^{n_1^m} \cdots b_k^{n_k^m}) c \in L$. By the closure properties of L , we have

$$a(b_1^{n_1^1} b_2^{n_2^1} \cdots b_k^{n_k^1}) \cdots (b_1^{n_1^m} \cdots b_k^{n_k^m}) c \in L.$$

An easy induction shows that $a(b_1^* \cdots b_k^*) \cdots (b_1^* \cdots b_k^*)^m c \in L$, that is, $a(b_1^* \cdots b_k^*)^m c \in L$, as required. $\square$

We can extend T to a functor by specifying its action on $\mathbf{PKA}^*$ morphisms. If $f : \mathcal{K} \rightarrow \mathcal{K}'$ is a homomorphism of partially additive Kleene algebras, define $Tf : T\mathcal{K} \rightarrow T\mathcal{K}'$ as $(Tf)(I) = \langle f[I] \rangle$, where $f[A] = \{f(a) \mid a \in A\}$.

Lemma 3.7. $\langle f[\langle A \rangle] \rangle = \langle f[A] \rangle$.

Proof. This is a straightforward adaption of the proof of Lemma 4 in Kozen [1990]. $\square$

Lemma 3.8. Tf is a Kleene algebra homomorphism.

Proof.

$$\begin{aligned} (Tf)(\langle 0 \rangle) &= \langle f[\langle 0 \rangle] \rangle \\ &= \langle f(0) \rangle && \text{by Lemma 3.7} \\ &= \langle 0 \rangle \end{aligned}$$

$$\begin{aligned}
(Tf)(\langle 1 \rangle) &= \langle f[\langle 1 \rangle] \rangle \\
&= \langle f[1] \rangle \quad \text{by Lemma 3.7} \\
&= \langle 1 \rangle.
\end{aligned}$$

Assume $I = \langle A \rangle$ and $J = \langle B \rangle$,

$$\begin{aligned}
(Tf)(I + J) &= \langle f[I + J] \rangle \\
&= \langle f[\langle A \cup B \rangle] \rangle \\
&= \langle f[A \cup B] \rangle \quad \text{by Lemma 3.7} \\
&= \langle f[A] \cup f[B] \rangle \\
&= \langle f[A] \rangle + \langle f[B] \rangle \\
&= \langle f[\langle A \rangle] \rangle + \langle f[\langle B \rangle] \rangle \quad \text{by Lemma 3.7} \\
&= \langle f[I] \rangle + \langle f[J] \rangle \\
&= (Tf)(I) + (Tf)(J)
\end{aligned}$$

$$\begin{aligned}
(Tf)(I \cdot J) &= \langle f[I \cdot J] \rangle \\
&= \langle f[\langle A \odot B \rangle] \rangle \\
&= \langle f[A \odot B] \rangle \quad \text{by Lemma 3.7} \\
&= \langle \{f(a \cdot b) \mid a \in A, b \in B\} \rangle \\
&= \langle \{f(a) \cdot f(b) \mid a \in A, b \in B\} \rangle \\
&= \langle \{x \cdot y \mid x \in f[A], y \in f[B]\} \rangle \\
&= \langle f[A] \odot f[B] \rangle \\
&= \langle f[A] \rangle \cdot \langle f[B] \rangle \\
&= \langle f[\langle A \rangle] \rangle \cdot \langle f[\langle B \rangle] \rangle \quad \text{by Lemma 3.7} \\
&= \langle f[I] \rangle \cdot \langle f[J] \rangle = (Tf)(I) \cdot (Tf)(J).
\end{aligned}$$

Assume $I = \langle \{a_1, \dots, a_k\} \rangle$,

$$\begin{aligned}
(Tf)(I^*) &= \langle f[I^*] \rangle \\
&= \langle f[\langle \{(a_1^* \cdots a_k^*)^* \} \rangle] \rangle \quad \text{by Lemma 3.7} \\
&= \langle f[\{(a_1^* \cdots a_k^*)^*\}] \rangle \\
&= \langle \{f((a_1^* \cdots a_k^*)^*)\} \rangle \\
&= \langle \{(f(a_1)^* \cdots f(a_k)^*)^*\} \rangle \\
&= (\langle \{f(a_1), \dots, f(a_k)\} \rangle)^* \\
&= (\langle f[\{a_1, \dots, a_k\}] \rangle)^* \\
&= (\langle f[\langle \{a_1, \dots, a_k\} \rangle] \rangle)^* \quad \text{by Lemma 3.7} \\
&= (\langle f[I] \rangle)^* = ((Tf)(I))^*.
\end{aligned}$$

□

It is straightforward to check that T is a functor.

Theorem 3.9. *The functor T is a left adjoint to the inclusion functor $P : \mathbf{KA}^* \rightarrow \mathbf{PKA}^*$, via the adjunction map $\varphi : \text{Hom}(T\mathcal{K}, \mathcal{K}') \rightarrow \text{Hom}(\mathcal{K}, PK')$ given by $(\varphi f)(x) = f(\langle x \rangle)$.*

Proof. To show that T is a left adjoint to P , we need to show that the adjunction map φ is a natural isomorphism between $\text{Hom}(T\mathcal{K}, \mathcal{K}')$ and $\text{Hom}(\mathcal{K}, PK')$.

We first show that (φf) is a partially additive Kleene algebra homomorphism. The verification is mostly straightforward.

$$\begin{aligned} (\varphi f)(0) &= f(\langle 0 \rangle) \\ &= \langle 0 \rangle, \end{aligned}$$

and similarly for 1.

$$\begin{aligned} (\varphi f)(x + y) &= f(\langle x + y \rangle) \\ &= f(\langle \{x, y\} \rangle) \\ &= f(\langle x \rangle + \langle y \rangle) \\ &= f(\langle x \rangle) + f(\langle y \rangle) \\ &= (\varphi f)(x) + (\varphi f)(y) \end{aligned}$$

$$\begin{aligned} (\varphi f)(x \cdot y) &= f(\langle x \cdot y \rangle) \\ &= f(\langle \{x\} \odot \{y\} \rangle) \\ &= f(\langle x \rangle \cdot \langle y \rangle) \\ &= f(\langle x \rangle) \cdot f(\langle y \rangle) \\ &= (\varphi f)(x) \cdot (\varphi f)(y) \end{aligned}$$

$$\begin{aligned} (\varphi f)(x^*) &= f(\langle x^* \rangle) \\ &= f(\langle (x^*)^* \rangle) \\ &= f(\langle x \rangle^*) \\ &= (f(\langle x \rangle))^* \\ &= ((\varphi f)(x))^*. \end{aligned}$$

We next show that φ is an isomorphism. The inverse of φ is given by the map ψ defined by $(\psi g)(I) = g(a_1) + \cdots + g(a_k)$ if $I = \langle a_1, \dots, a_k \rangle$. We check that this function is well defined, that is, if $I = \langle a_1, \dots, a_k \rangle$ and $I = \langle b_1, \dots, b_l \rangle$, then $g(a_1) + \cdots + g(a_k) = g(b_1) + \cdots + g(b_l)$. First, since $g : \mathcal{K} \rightarrow PK'$ and since P is the forgetful functor from $\mathbf{KA}^*$, the operation $+$ is total on the range of g . Note that from Lemma 3.7, we have that $\langle g[I] \rangle = \langle g(a_1), \dots, g(a_k) \rangle = \langle g(b_1), \dots, g(b_l) \rangle \subseteq \langle g(b_1) + \cdots + g(b_l) \rangle = (g(b_1) + \cdots + g(b_l))^\leq$. Therefore, $g(a_i) \leq g(b_1) + \cdots + g(b_l)$ for all i , and therefore $g(a_1) + \cdots + g(a_k) \leq g(b_1) + \cdots + g(b_l)$. A symmetric argument shows that $g(b_1) + \cdots + g(b_l) \leq g(a_1) + \cdots + g(a_k)$, establishing the result. We check that (ψg) is a Kleene algebra homomorphism.

$$\begin{aligned} (\psi g)(\langle 0 \rangle) &= g(0) \\ &= 0 \end{aligned}$$

and similarly for 1.

$$\begin{aligned}
(\psi g)(\langle A \rangle + \langle B \rangle) &= (\psi g)(\langle A \cup B \rangle) \\
&= \sum g[A \cup B] \\
&= \sum g[A] \cup g[B] \\
&= \sum g[A] + \sum g[B] \\
&= (\psi g)(\langle A \rangle) + (\psi g)(\langle B \rangle)
\end{aligned}$$

$$\begin{aligned}
(\psi g)(\langle A \rangle \cdot \langle B \rangle) &= (\psi g)(\langle A \odot B \rangle) \\
&= \sum g[A \odot B] \\
&= \sum g[A] \odot g[B] \\
&= \sum g[A] \cdot \sum g[B] \\
&= (\psi g)(\langle A \rangle) \cdot (\psi g)(\langle B \rangle)
\end{aligned}$$

$$\begin{aligned}
(\psi g)(\langle \{a_1, \dots, a_k\} \rangle^*) &= (\psi g)(\langle (a_1^* \cdots a_k^*)^* \rangle) \\
&= g((a_1^* \cdots a_k^*)^*) \\
&= (g(a_1)^* \cdots g(a_k)^*)^* \\
&= (g(a_1) + \cdots + g(a_k))^* \\
&= (\sum g[\{a_1, \dots, a_k\}])^* \\
&= ((\psi g)(\langle a_1, \dots, a_k \rangle))^*.
\end{aligned}$$

To see that φ and ψ are inverses, first let $g : \mathcal{K} \rightarrow P\mathcal{K}'$, and note that

$$\begin{aligned}
(\varphi \circ \psi)(g)(x) &= (\varphi(\psi(g)))(x) \\
&= (\psi g)(\langle x \rangle) \\
&= g(x).
\end{aligned}$$

Similarly, let $f : T\mathcal{K} \rightarrow \mathcal{K}'$, and note that

$$\begin{aligned}
(\psi \circ \varphi)(f)(\langle \{a_1, \dots, a_k\} \rangle) &= (\psi(\varphi(f)))(\langle \{a_1, \dots, a_k\} \rangle) \\
&= (\varphi f)(a_1) + \cdots + (\varphi f)(a_k) \\
&= f(\langle a_1 \rangle) + \cdots + f(\langle a_k \rangle) \\
&= f(\langle a_1 \rangle + \cdots + \langle a_k \rangle) \\
&= f(\langle a_1, \dots, a_k \rangle).
\end{aligned}$$

To establish the naturality of the adjunction map φ , we need to show the commutativity of two diagrams. (See the proof of Theorem 3.2.) First, let $h : \mathcal{K}' \rightarrow \mathcal{K}''$, let $f : T\mathcal{K} \rightarrow \mathcal{K}'$, and let $x \in \mathcal{K}$:

$$\begin{aligned}
(\text{Hom}(\mathcal{K}, Ph) \circ \varphi)(f)(x) &= (\text{Hom}(\mathcal{K}, Ph)(\varphi(f)))(x) \\
&= ((Ph) \circ (\varphi f))(x) \\
&= ((Ph)((\varphi f)(x))) \\
&= h(f(\langle x \rangle))
\end{aligned}$$

(since $Ph = h$) and

$$\begin{aligned}
(\varphi \circ \text{Hom}(TK, h))(f)(x) &= (\varphi(\text{Hom}(TK, h)(f)))(x) \\
&= (\varphi(h \circ f))(x) \\
&= (h \circ f)(\langle x \rangle) \\
&= h(f(\langle x \rangle)).
\end{aligned}$$

Second, let $h : \mathcal{K}'' \rightarrow \mathcal{K}'$, let $f : TK' \rightarrow \mathcal{K}$, and let $x \in \mathcal{K}''$:

$$\begin{aligned}
(\text{Hom}(h, PK) \circ \varphi)(f)(x) &= (\text{Hom}(h, PK)(\varphi f))(x) \\
&= ((\varphi f) \circ h)(x) \\
&= (\varphi f)(h(x)) \\
&= f(\langle h(x) \rangle)
\end{aligned}$$

and

$$\begin{aligned}
(\varphi \circ \text{Hom}(Th, \mathcal{K}))(f)(x) &= (\varphi(\text{Hom}(Th, \mathcal{K})(f)))(x) \\
&= (\varphi(f \circ (Th)))(x) \\
&= (f \circ (Th))(\langle x \rangle) \\
&= f((Th)(\langle x \rangle)) \\
&= f(\langle h(x) \rangle).
\end{aligned}$$

□

4 Relationship with Partially Additive Idempotent Semirings

An algebraic structure that is quite similar to partially additive Kleene algebras was studied by Arbib and Manes [1986], who introduced a form of partially additive semiring with an infinitary sum operation.¹ At the basis of this structure is the notion of a *partially additive monoid*. Roughly speaking, a partially additive monoid is like a monoid (that is, a set with a single associate operation and an identity element), except that the operation is only partially defined, and is infinitary. In this section, we show the relationship between partially additive Kleene algebras and partially additive idempotent semirings.

To make this precise, we need some terminology. Let $\mathcal{M}$ be a fixed set. An $\mathcal{I}$ -indexed family in $\mathcal{M}$ is a function $x : \mathcal{I} \rightarrow \mathcal{M}$, written $(x_i \mid i \in \mathcal{I})$. We usually write x_i rather than $x(i)$. The (necessarily unique) $\emptyset$ -indexed family is called the *empty family*. A family $(y_j \mid j \in \mathcal{J})$ is said to be a *subfamily* of $(x_i \mid i \in \mathcal{I})$ if $\mathcal{J} \subseteq \mathcal{I}$ and $y_j = x_j$ for all $j \in \mathcal{J}$. A family $(x_i \mid i \in \mathcal{I})$ is *countable* if $\mathcal{I}$ is a countable set. Given an index set $\mathcal{I}$, the family $(\mathcal{I}_j \mid j \in \mathcal{J})$ is a *partition* of $\mathcal{I}$ if $\mathcal{I}_j \cap \mathcal{I}_k = \emptyset$ whenever $j \neq k$, and $\mathcal{I} = \cup_{j \in \mathcal{J}} \mathcal{I}_j$. Note that we allow an element $\mathcal{I}_j$ of the partition to be empty.

A *partially additive monoid* is a pair $(\mathcal{M}, \sum)$ where $\mathcal{M}$ is a nonempty set and $\sum$ is a partial function mapping countable families in $\mathcal{M}$ to elements of $\mathcal{M}$ (the family $(x_i \mid i \in \mathcal{I})$ is *summable* if $\sum(x_i \mid i \in \mathcal{I})$ is defined) subject to the following three conditions:

¹Strictly speaking, Arbib and Manes define the notion of a *partially additive category*. A partially additive semiring is a one-object partially additive category.

- (1) *Partition-associativity axiom:* If $(x_i \mid i \in \mathcal{I})$ is a countable family and $(\mathcal{I}_j \mid j \in \mathcal{J})$ is a partition of $\mathcal{I}$ with $\mathcal{J}$ countable, then $(x_i \mid i \in \mathcal{I})$ is summable if and only if the case both that for every $j \in \mathcal{J}$, $(x_i \mid i \in \mathcal{I}_j)$ is summable, and $(\sum(x_i \mid i \in \mathcal{I}_j) \mid j \in \mathcal{J})$ is summable. In that case, $\sum(x_i \mid i \in \mathcal{I}) = \sum(\sum(x_i \mid i \in \mathcal{I}_j) \mid j \in \mathcal{J})$;
- (2) *Unary sum axiom:* Any family $(x_i \mid i \in \mathcal{I})$ where $|\mathcal{I}| = 1$ is summable, and $\sum(x_i \mid i \in \mathcal{I}) = x_j$ if $\mathcal{I} = \{j\}$;
- (3) *Limit axiom:* If $(x_i \mid i \in \mathcal{I})$ is a countable family, and if the subfamily $(x_i \mid i \in F)$ is summable for every finite set $F \subseteq \mathcal{I}$, then $(x_i \mid i \in \mathcal{I})$ is summable.

One consequence of the partition-associativity axiom is that every partially additive monoid is commutative: the order of the elements of the family being summed does not affect the sum. Moreover, in every partially additive monoid there exists a special element that acts as the identity under sums. More precisely, for a partially additive monoid $\mathcal{M}$, let $! : \emptyset \longrightarrow \mathcal{M}$ be the empty family in $\mathcal{M}$, which is summable by an application of the partition-associativity axiom. Let 0 be $\sum!$. That this 0 acts as the identity is captured by the following easily-proved fact: if $(x_i \mid i \in \mathcal{I})$ is a summable family in $\mathcal{M}$, $\mathcal{J}$ a countable set disjoint from $\mathcal{I}$, and $x_i = 0$ for $i \in \mathcal{J}$, then $(x_i \mid i \in \mathcal{I} \cup \mathcal{J})$ is summable, and $\sum(x_i \mid i \in \mathcal{I} \cup \mathcal{J}) = \sum(x_i \mid i \in \mathcal{I})$.

A *partially additive idempotent semiring* is a structure $(\mathcal{S}, \sum, \cdot, 1)$ where

- (1) $(\mathcal{S}, \sum)$ is a partially additive monoid that is idempotent: if $\mathcal{I}$ is a countable set and we form the family $(x_i \mid i \in \mathcal{I})$ with $x_i = x$ for all $i \in \mathcal{I}$, then $(x_i \mid i \in \mathcal{I})$ is summable, and $\sum(x_i \mid i \in \mathcal{I}) = x$;
- (2) $(\mathcal{S}, \cdot, 1)$ is a monoid;
- (3) If $(x_i \mid i \in \mathcal{I})$ is summable, then $(y \cdot x_i \mid i \in \mathcal{I})$ and $(x_i \cdot z \mid i \in \mathcal{I})$ are summable, and $y(\sum(x_i \mid i \in \mathcal{I})) = \sum(y \cdot x_i \mid i \in \mathcal{I})$ and $(\sum(x_i \mid i \in \mathcal{I}))z = \sum(x_i \cdot z \mid i \in \mathcal{I})$;
- (4) $a \cdot 0 = 0 \cdot a = 0$.

Partially additive idempotent semirings form a category **PS**, with morphisms being homomorphisms that preserve the sum of summable countable families. (These homomorphisms are sometimes called ω -continuous.)

In the examples treated in the literature, the infinitary sum is used almost exclusively to define the $*$ operation, by taking $b^* = \sum(b^n \mid n \in \mathbb{N})$, where $b^0 = 1$ and $b^{n+1} = bb^n$. By countable distributivity, we have $ab^*c = \sum(ab^n c \mid n \in \mathbb{N})$, and therefore every partially additive idempotent semiring is a star-continuous partially additive Kleene algebra. This gives us a forgetful functor $U : \mathbf{PS} \rightarrow \mathbf{PKA}^*$.

There is a way to universal embed a star-continuous partially additive Kleene algebra into a partially additive idempotent semiring, by essentially extending the $+$ operation to an infinitary summation. This takes the form of a functor C from $\mathbf{PKA}^*$ to $\mathbf{PS}$. It is again a form of ideal completion, of the kind already seen in Section 3. We use the same definition of star-ideals. This time, however, we take CK to be the set of *countably generated* star-ideals of $\mathcal{K}$. We define the partially additive idempotent semiring operations as follows. Most of the operations are defined as

we did for the functor T in Section 3. If I and J are countably generated star-ideals, generated respectively by the countable sets A and B , define

$$\begin{aligned} I \cdot J &= \langle A \odot B \rangle \\ 1 &= \langle 1 \rangle \\ 0 &= \langle 0 \rangle. \end{aligned}$$

To define the infinitary summation $\sum$, let $(I_i \mid i \in \mathcal{I})$ be a countable family of countably generated star-ideals of $\mathcal{K}$, where I_i is generated by the countable set A_i . Say that the family $(I_i \mid i \in \mathcal{I})$ is summable if for all finite subsets $\{a_1, \dots, a_k\} \subseteq \bigcup_{i \in \mathcal{I}} A_i$, the finite sum $a_1 + \dots + a_k$ is defined in $\mathcal{K}$, for all possible ways of associating the binary $+$ operations in $a_1 + \dots + a_k$; if $(I_i \mid i \in \mathcal{I})$ is summable, define $\sum(I_i \mid i \in \mathcal{I}) = \langle \bigcup_{i \in \mathcal{I}} A_i \rangle$. Again, we can check that these operations are well-defined, by showing that they give the same star-ideals irrespectively of the choice of countable generating sets. (The details are as in Lemma 3.5.)

It is straightforward to check that CK is a partially additive idempotent semiring. We can extend C to a functor by specifying its action on $\mathbf{PKA}^*$ morphisms. If $f : \mathcal{K} \rightarrow \mathcal{K}'$ is a homomorphism of partially additive Kleene algebras, define $Cf : CK \rightarrow CK'$ as $(Cf)(I) = \langle f[I] \rangle$, where $f[A] = \{f(a) \mid a \in A\}$.

Lemma 4.1. *Cf is a partially additive idempotent semiring homomorphism.*

Proof. This proof is similar to that of Lemma 3.8. The only case that is different (aside from the fact that there is no case for $*$, since there is no such operation in partially additive idempotent semirings) is the countable sum. Let $(I_i \mid i \in \mathcal{I})$ be a countable family of countably generated star-ideals, where $I_i = \langle A_i \rangle$. Assume $(I_i \mid i \in \mathcal{I})$ is summable.

$$\begin{aligned} (Cf)(\sum(I_i \mid i \in \mathcal{I})) &= \langle f[\sum(I_i \mid i \in \mathcal{I})] \rangle \\ &= \langle f[\langle \bigcup_{i \in \mathcal{I}} A_i \rangle] \rangle \\ &= \langle f[\bigcup_{i \in \mathcal{I}} A_i] \rangle \quad \text{by Lemma 3.7} \\ &= \langle \bigcup_{i \in \mathcal{I}} f[A_i] \rangle \\ &= \sum(\langle f[A_i] \rangle \mid i \in \mathcal{I}) \\ &= \sum(\langle f[\langle A_i \rangle] \rangle \mid i \in \mathcal{I}) \quad \text{by Lemma 3.7} \\ &= \sum(\langle f[I_i] \rangle \mid i \in \mathcal{I}) \\ &= \sum((Cf)(I_i) \mid i \in \mathcal{I}). \end{aligned}$$

□

It is straightforward to check that C is a functor.

Theorem 4.2. *The functor C is a left adjoint of the inclusion functor $U : \mathbf{PS} \rightarrow \mathbf{PKA}^*$, via the adjunction map $\varphi : \text{Hom}(CK, S) \rightarrow \text{Hom}(\mathcal{K}, US)$ given by $(\varphi f)(x) = f(\langle x \rangle)$.*

Proof. To show that C is a left adjoint to U , we need to show that the adjunction map φ is a natural isomorphism between $\text{Hom}(CK, \mathcal{S})$ and $\text{Hom}(K, US)$.

We first show that (φf) is a partially additive Kleene algebra homomorphism. The verification is mostly straightforward.

$$\begin{aligned}(\varphi f)(0) &= f(\langle 0 \rangle) \\ &= \langle 0 \rangle,\end{aligned}$$

and similarly for 1.

$$\begin{aligned}(\varphi f)(x + y) &= f(\langle x + y \rangle) \\ &= f(\langle \{x, y\} \rangle) \\ &= f(\langle x \rangle + \langle y \rangle) \\ &= f(\langle x \rangle) + f(\langle y \rangle) \\ &= (\varphi f)(x) + (\varphi f)(y)\end{aligned}$$

$$\begin{aligned}(\varphi f)(x \cdot y) &= f(\langle x \cdot y \rangle) \\ &= f(\langle \{x\} \odot \{y\} \rangle) \\ &= f(\langle x \rangle \cdot \langle y \rangle) \\ &= f(\langle x \rangle) \cdot f(\langle y \rangle) \\ &= (\varphi f)(x) \cdot (\varphi f)(y).\end{aligned}$$

For x^* , note that $\langle x^* \rangle = \langle \bigcup_{n \geq 0} \{x^n\} \rangle$: since $x^n \leq x^*$ for all $n \geq 0$, $\bigcup_{n \geq 0} \{x^n\} \subseteq \langle x^* \rangle$ by the closure properties of star-ideals, and thus $\langle \bigcup_{n \geq 0} \{x^n\} \rangle \subseteq \langle x^* \rangle$; conversely, we have $x^n \in \langle \bigcup_{n \geq 0} \{x^n\} \rangle$ for all $n \geq 0$, and again by the closure properties of star-ideals, $x^* \in \langle \bigcup_{n \geq 0} \{x^n\} \rangle$, and thus $\langle x^* \rangle \subseteq \langle \bigcup_{n \geq 0} \{x^n\} \rangle$.

$$\begin{aligned}(\varphi f)(x^*) &= f(\langle x^* \rangle) \\ &= f(\langle \bigcup_{n \geq 0} \{x^n\} \rangle) \\ &= f(\sum(\langle x^n \rangle \mid n \geq 0)) \\ &= f(\sum(\langle x \rangle^n \mid n \geq 0)) \\ &= \sum(f(\langle x \rangle)^n \mid n \geq 0) \\ &= \sum((\varphi f)(x)^n \mid n \geq 0) \\ &= ((\varphi f)(x))^* \quad \text{by definition of } *.\end{aligned}$$

We next show that φ is an isomorphism. The inverse of φ is given by the map ψ defined by $(\psi g)(I) = \sum(g(a) \mid a \in A)$ if $I = \langle A \rangle$. We check that this function is well defined, that is, if $I = \langle A \rangle$ and $I = \langle B \rangle$, then $\sum(g(a) \mid a \in A) = \sum(g(b) \mid b \in B)$. The argument is essentially the same as that in the proof of Theorem 3.9. We repeat the argument here for convenience. Note that from Lemma 3.7, we have that $\langle g[I] \rangle = \langle g[A] \rangle = \langle g[B] \rangle \subseteq \langle \sum(g(b) \mid b \in B) \rangle = \langle \sum(g(b) \mid$

$b \in B)) \leq$. Therefore, $g(a) \leq \sum(g(b) \mid b \in B)$ for all $a \in A$, and therefore $\sum(g(a) \mid a \in A) \leq \sum(g(b) \mid b \in B)$. A symmetric argument shows that $\sum(g(b) \mid b \in B) \leq \sum(g(a) \mid a \in A)$, establishing the result. We check that (ψg) is a partially additive idempotent semiring homomorphism.

$$\begin{aligned} (\psi g)(\langle 0 \rangle) &= g(0) \\ &= 0 \end{aligned}$$

and similarly for 1.

$$\begin{aligned} (\psi g)\left(\sum(\langle A_i \rangle \mid i \in \mathcal{I})\right) &= (\psi g)\left(\left\langle \bigcup_{i \in \mathcal{I}} A_i \right\rangle\right) \\ &= \sum(g(a) \mid a \in \bigcup_{i \in \mathcal{I}} A_i) \\ &= \sum\left(\sum(g(a) \mid a \in A_i) \mid i \in \mathcal{I}\right) \\ &= \sum((\psi g)(\langle A_i \rangle) \mid i \in \mathcal{I}) \end{aligned}$$

$$\begin{aligned} (\psi g)(\langle A \rangle \cdot \langle B \rangle) &= (\psi g)(\langle A \odot B \rangle) \\ &= \sum g[A \odot B] \\ &= \sum g[A] \odot g[B] \\ &= \sum g[A] \cdot \sum g[B] \\ &= (\psi g)(\langle A \rangle) \cdot (\psi g)(\langle B \rangle). \end{aligned}$$

To see that φ and ψ are inverses, first let $g : \mathcal{K} \rightarrow \mathcal{US}$, and note that

$$\begin{aligned} (\varphi \circ \psi)(g)(x) &= (\varphi(\psi(g)))(x) \\ &= (\psi g)(\langle x \rangle) \\ &= g(x). \end{aligned}$$

Similarly, let $f : C\mathcal{K} \rightarrow \mathcal{S}$, and note that

$$\begin{aligned} (\psi \circ \varphi)(f)(\langle A \rangle) &= (\psi(\varphi(f)))(\langle A \rangle) \\ &= \sum((\varphi f)(a) \mid a \in A) \\ &= \sum(f(\langle a \rangle) \mid a \in A) \\ &= f\left(\sum(\langle a \rangle \mid a \in A)\right) \\ &= f(\langle A \rangle). \end{aligned}$$

To establish the naturality of the adjunction map φ , we need to show the commutativity of two diagrams. (See the proof of Theorem 3.2.) First, let $h : \mathcal{S} \rightarrow \mathcal{S}'$, let $f : C\mathcal{K} \rightarrow \mathcal{S}$, and let $x \in \mathcal{K}$:

$$\begin{aligned} (\text{Hom}(\mathcal{K}, Uh) \circ \varphi)(f)(x) &= (\text{Hom}(\mathcal{K}, Uh)(\varphi(f)))(x) \\ &= ((Uh) \circ (\varphi f))(x) \\ &= ((Uh)((\varphi f)(x))) \\ &= h(f(\langle x \rangle)) \end{aligned}$$

(since $Uh = h$) and

$$\begin{aligned}
(\varphi \circ \text{Hom}(CK, h))(f)(x) &= (\varphi(\text{Hom}(CK, h)(f)))(x) \\
&= (\varphi(h \circ f))(x) \\
&= (h \circ f)(\langle x \rangle) \\
&= h(f(\langle x \rangle)).
\end{aligned}$$

Second, let $h : \mathcal{K}' \rightarrow \mathcal{K}$, let $f : CK \rightarrow \mathcal{S}$, and let $x \in \mathcal{K}'$:

$$\begin{aligned}
(\text{Hom}(h, US) \circ \varphi)(f)(x) &= (\text{Hom}(h, US)(\varphi f))(x) \\
&= ((\varphi f) \circ h)(x) \\
&= (\varphi f)(h(x)) \\
&= f(\langle h(x) \rangle)
\end{aligned}$$

and

$$\begin{aligned}
(\varphi \circ \text{Hom}(Ch, \mathcal{S}))(f)(x) &= (\varphi(\text{Hom}(Ch, \mathcal{S})(f)))(x) \\
&= (\varphi(f \circ (Ch)))(x) \\
&= (f \circ (Ch))(\langle x \rangle) \\
&= f((Ch)(\langle x \rangle)) \\
&= f(\langle h(x) \rangle).
\end{aligned}$$

□

5 Relationship with Closed Semirings

Closed semirings [Aho, Hopcroft, and Ullman 1975; Mehlhorn 1984] are often used as alternatives to Kleene algebras to treat iteration by defining it via an infinite sum operation. In fact, a closed semiring is simply a partially additive idempotent semiring where the $\sum$ operator is always defined, that is, where every countable family is summable. In this section, we show that there are two ways to universally map star-continuous partially additive Kleene algebras to closed semirings, using the constructions of the previous two sections; these nevertheless yield isomorphic closed semirings.

In Section 3, we showed that star-continuous partially additive Kleene algebras can be completed to star-continuous Kleene algebras, via the functor T . Kozen [1990] showed that every star-continuous Kleene algebra can be universally embedded into a closed semiring, via a functor $C' : \mathbf{KA}^* \rightarrow \mathbf{CS}$ (where $\mathbf{CS}$ is the category of closed semirings) and that C' is a left adjoint to the inclusion functor $U' : \mathbf{CS} \rightarrow \mathbf{KA}^*$. By composing the functors, we get the adjunctions

$$\mathbf{PKA}^* \begin{array}{c} \xrightarrow{T} \\ \xleftarrow{P} \end{array} \mathbf{KA}^* \begin{array}{c} \xrightarrow{C'} \\ \xleftarrow{U'} \end{array} \mathbf{CS}.$$

Thus, the composite functor $C' \circ T$ is a left adjoint to the forgetful functor $\mathbf{CS} \rightarrow \mathbf{PKA}^*$.

In Section 4 we showed that every star-continuous partially additive Kleene algebra can be embedded into a partially additive idempotent semirings, via the functor $C : \mathbf{PKA}^* \rightarrow \mathbf{PS}$. It

is straightforward to show that partially additive idempotent semirings can be completed to closed semirings in a universal way; this construction is quite similar to the completion of partially additive Kleene algebras. This construction is essentially the one described by Manes [1987]. If $\mathcal{S}$ is a partially additive idempotent semiring, define an *ideal* of $\mathcal{S}$ to be a subset $A \subseteq \mathcal{S}$ satisfying:

- (1) A is nonempty;
- (2) A is closed under $\sum$ (for summable families);
- (3) A is closed downward under $\leq$.

We use the same notation as for star-ideals, and write $\langle A \rangle$ for the smallest ideal generated by the set A . The analogue of results for star-ideals carry over (for instance, Lemmas 3.3 and 3.4). If $\mathcal{S}$ is a partially additive idempotent semiring, let $T'\mathcal{S}$ be the set of ideals of $\mathcal{S}$. We can impose a closed semiring structure on $T'\mathcal{S}$ as follows. If I and J are ideals, generated respectively by A and B , define

$$\begin{aligned}\sum (I_i \mid i \in \mathcal{I}) &= \langle \bigcup_{i \in \mathcal{I}} A_i \rangle \\ I \cdot J &= \langle A \odot B \rangle \\ 1 &= \langle 1 \rangle \\ 0 &= \langle 0 \rangle.\end{aligned}$$

As in Section 3, we need to check that these operations are well-defined; the analogue of Lemma 3.5 can be seen to hold. As with T , we can extend T' to a functor by specifying its action on $\mathbf{PS}$ morphisms. If $f : \mathcal{S} \rightarrow \mathcal{S}'$ is a homomorphism of partially additive idempotent semirings, define $T'f : T'\mathcal{S} \rightarrow T'\mathcal{S}'$ as $(T'f)(I) = \langle f[I] \rangle$, where $f[A] = \{f(a) \mid a \in A\}$. It is straightforward to check that $T'f$ is a closed semiring homomorphism (see Lemma 3.8). This gives a functor T' from the category $\mathbf{PS}$ to the category $\mathbf{CS}$, left adjoint to the obvious inclusion functor $P' : \mathbf{CS} \rightarrow \mathbf{PS}$.

Theorem 5.1. *The functor T' is a left adjoint to the inclusion functor $P' : \mathbf{CS} \rightarrow \mathbf{PS}$, via the adjunction map $\varphi : \text{Hom}(T'\mathcal{S}, \mathcal{S}') \rightarrow \text{Hom}(\mathcal{S}, P'\mathcal{S}')$ given by $(\varphi f)(x) = f(\langle x \rangle)$.*

Proof. To show that T' is a left adjoint to P' , we need to show that the adjunction map φ is a natural isomorphism between $\text{Hom}(T'\mathcal{S}, \mathcal{S}')$ and $\text{Hom}(\mathcal{S}, P'\mathcal{S}')$. The proof is a straightforward adaptation of the proof of Theorem 3.9. $\square$

With this result, we get the adjunctions

$$\mathbf{PKA}^* \begin{array}{c} \xrightarrow{C} \\ \xleftarrow{U} \end{array} \mathbf{PS} \begin{array}{c} \xrightarrow{T'} \\ \xleftarrow{P'} \end{array} \mathbf{CS}.$$

The composite functor $T' \circ C$ is a left adjoint to the forgetful functor $\mathbf{CS} \rightarrow \mathbf{PKA}^*$. It is a well-known result that any two functors that are left adjoint to the same functor are naturally isomorphic [Mac Lane 1971, p.85, Corollary 1]. Thus, $C' \circ T$ and $T' \circ C$ are naturally isomorphic functors. Among other things, this implies that if $\mathcal{K}$ is a star-continuous partially additive Kleene algebra, then $C'(TK)$ and $T'(CK)$ are isomorphic closed semirings.

6 Conclusion

In this paper, we introduced the notion of a partially additive Kleene algebra, and exhibited categorical connections between the category $\mathbf{PKA}^*$ of star-continuous partially additive Kleene algebras and various other algebraic structures. We obtain the following commutative diagram of adjunctions:

$$\begin{array}{ccccc}
 \mathbf{PKA} & \xrightleftharpoons[S]{K} & \mathbf{PKA}^* & \xrightleftharpoons[U]{C} & \mathbf{PS} \\
 & & \uparrow P \quad \downarrow T & & \uparrow P' \quad \downarrow T' \\
 & & \mathbf{KA}^* & \xrightleftharpoons[U']{C'} & \mathbf{CS}
 \end{array}$$

These relationships provide some motivation for our definition of partially additive Kleene algebras, by showing that partially additive Kleene algebras fit naturally within existing algebraic structures.

One question remains open: is there is a universal way to complete an arbitrary partially additive Kleene algebra into a total Kleene algebra? This asks for a left adjoint to the forgetful functor $\mathbf{KA} \rightarrow \mathbf{PKA}$, that collapses to the functor $T : \mathbf{PKA}^* \rightarrow \mathbf{KA}^*$ when restricted to $\mathbf{PKA}^*$. Note that the construction in this paper only works for star-continuous partially additive Kleene algebras. More precisely, the proof of Theorem 3.9 requires showing that the adjunction φ is an isomorphism; but the well-definedness of the inverse ψ relies on the fact that $\langle x \rangle \subseteq \{x\}^\leq$, which need not hold in non-star-continuous partially additive Kleene algebras. For instance, if y is an upper bound for all x^n but not x^* , then $\langle y \rangle$ will contain x^* but $\{y\}^\leq$ will not; in other words, $\langle y \rangle \not\subseteq \{y\}^\leq$.

Acknowledgments. Thanks to Dexter Kozen for an enlightening comment pointing out the necessity of star-continuity. Thanks as well to Hubie Chen for commenting on a draft of this paper.

References

- Aho, A. V., J. E. Hopcroft, and J. D. Ullman (1975). *The Design and Analysis of Computer Algorithms*. Addison Wesley.
- Conway, J. H. (1971). *Regular Algebra and Finite Machines*. Chapman and Hall.
- Kozen, D. (1990). On Kleene algebras and closed semirings. In *Proc. 15th International Symposium on the Mathematical Foundations of Computer Science (MFCS'90)*, Volume 452 of *Lecture Notes in Computer Science*, pp. 26–47. Springer-Verlag.
- Kozen, D. (1994). A completeness theorem for Kleene algebras and the algebra of regular events. *Information and Computation* 110, 366–390.
- Kozen, D. (1998). Typed Kleene algebra. Technical Report 98-1669, Computer Science Department, Cornell University.
- Kozen, D. (2003). Kleene algebra and dataflow analysis. In *Proceedings of the 7th International Seminar on Relational Methods in Computer Science*.

- Mac Lane, S. (1971). *Categories for the Working Mathematician*. Number 5 in Graduate Texts in Mathematics. Springer-Verlag.
- Manes, E. (1987). Assertion categories. In *Proc. 3rd Workshop on the Mathematical Foundations of Programming Semantics*, Volume 298 of *Lecture Notes in Computer Science*, pp. 85–120. Springer-Verlag.
- Manes, E. and M. Arbib (1986). *Algebraic Approaches to Program Semantics*. Springer-Verlag.
- Mehlhorn, K. (1984). *Data Structures and Algorithms 2: Graph Algorithms and NP-Completeness*. EATCS Monographs in Theoretical Computer Science. Springer-Verlag.
- Tarski, A. (1955). A lattice-theoretical fixpoint theorem and its applications. *Pacific Journal of Mathematics* 5, 285–309.